

GFastGene® Restriction Enzyme BtsC I



Cat.# FG-BtsCi Size 2.000 units Conc. 20 units/µl

Store at -20°C

Supplied with: 10X FastGene® Buffer IV (FG-REB4) 10X FastGene® FastCut Buffer (FG-REBHF) 6X DNA Loading Buffer Sterile water

Recognition site



For Research Use Only. Not for use in diagnostic procedures.

Dilution buffer:

FastGene® Diluent B

Heat Inactivation

BtsC I can be inactivated at 80°C for 20 min.

Methylation sensitivity

dam methylation: Not sensitive *dcm* methylation: Not sensitive CpG methylation: Not sensitive

Prolonged incubation

A minimum amount of enzyme required to digest 1 μg substrate DNA for 16 hr; 0.5 U.

Relative activity in FastGene® Buffers

FastGene [®] Buffe	rl: 75%
FastGene [®] Buffe	r II: 100%
FastGene [®] Buffe	r III: 100%
FastGene® Buffe	r IV: 100%
FastGene® FastC	Cut Buffer: 100%

Note

It is an isoschizomer of BstF5 I. It is not affected by *dam, dcm,* or mammalian CpG methylation.

Source: Bacillus thermosphaericus

Reaction conditions

1X FastGene[®] Buffer IV 50°C 1X FastGene[®] FastCut Buffer, 50°C

FastGene® FastCut Buffer

FastGene® restriction enzyme can cut substrate DNA in 5-15 with FastGene® FastCut Buffer.

1X FastGene® Buffer IV

20 mM Tris-acetate (pH 7.9 at 25°C) 50 mM potassium acetate 10 mM magnesium acetate 100 μg/ml BSA

Unit definition

One unit is defined as the amount of enzyme required for complete digestion of 1 μ g bacteriophage λ at 50°C for 1 hr in 50 μ l reaction mixtures.

Quality control

- Unit definition assay
- Overdigestion assay
- Endonuclease assay
- Extreme pure assay

Standard reaction condition

 Normal protocol

Component	Final Conc.	Volume
Substrate DNA	1 µg	Xμl
10X FastGene [®] Buffer IV	1 X	5 µl
BtsC I	20 unit	1 µl
Sterile water		up to 50 µl
, Incubate at 50℃ for 1 br		

→ Incubate at 50°C for 1 h

- Fast protocol

Component	Final Conc.	Volume
Substrate DNA	1 µg	X µl
10X FastGene [®] FastCut Buffer	1 X	5 µl
BtsC I	20 unit	1 µl
Sterile water		up to 50 µl

 \rightarrow Incubate at 50°C for 15 min

% We recommend 5-10 units of enzyme per μg DNA and 10-20 units for genomic DNA in a 1 h digest.