

G Fast Gene **Restriction Enzyme**



Conc

Cat.# FG-Avrll

Avr II

Size 100 units

4 units/ul

Store at -20°C

Supplied with: 10X FastGene® Buffer IV (FG-REB4) 10X FastGene® FastCut Buffer (FG-REBHF) 6X DNA Loading Buffer Sterile water

Recognition site



For Research Use Only. Not for use in diagnostic procedures.

ISO9001

Dilution buffer:

FastGene® Diluent B

Heat Inactivation

Avr II can be inactivated at 80°C for 20 min.

Methylation sensitivity

dam methylation: Not sensitive dcm methylation: Not sensitive CpG methylation: Not sensitive

Prolonged incubation

A minimum amount of enzyme required to digest 1 µg substrate DNA for 16 hr; 0.13 U.

Relative activity in FastGene® Buffers

FastGene®	Buffer I:	100%
FastGene®	Buffer II:	50%
FastGene®	Buffer III:	50%
FastGene®	Buffer IV:	100%
FastGene®	FastCut Buffer:	100%

Note

It is not affected by dam, dcm, or mammalian CpG methylation.

Source: Anabaena variabilis UW

Reaction conditions

1X FastGene® Buffer IV 37°C 1X FastGene® FastCut Buffer, 37°C

FastGene[®] FastCut Buffer

FastGene® restriction enzyme can cut substrate DNA in 5-15 with FastGene® FastCut Buffer.

1X FastGene® Buffer IV

20 mM Tris-acetate (pH 7.9 at 25°C) 50 mM potassium acetate 10 mM magnesium acetate 100 µg/ml BSA

Unit definition

One unit is defined as the amount of enzyme required for complete digestion of 1 μ g bacteriophage λ (Hind III digestion) at 37°C for 1 hr in 50 µl reaction mixtures.

Quality control

- Unit definition assay
- Overdigestion assay
- Endonuclease assay
- Extreme pure assay

Standard reaction condition

 Normal 	protocol	
----------------------------	----------	--

Component	Final Conc.	Volume
Substrate DNA	1 µg	X µl
10X FastGene [®] Buffer IV	1 X	5 µl
Avr II	4 unit	1 µl
Sterile water		up to 50 µl
\rightarrow Incubate at 37°C for 1 hr		

Incubate at 3/°C for 1 h

-	Fast	protocol	

Component	Final Conc.	Volume
Substrate DNA	1 µg	X µl
10X FastGene® FastCut Buffer	1 X	5 µl
Avr II	4 unit	1 µl
Sterile water		up to 50 µl
1 h		

→ Incubate at 37°C for 15 min

※ We recommend 5-10 units of enzyme per μg DNA and 10-20 units for genomic DNA in a 1 h digest.