



## Restriction Enzyme

### BstY I




Cat.# **FG-BstYI**      Size **2,000 units**      Conc. **10 units/μl**

Store at **-20°C**

**Supplied with:** 10X FastGene® Buffer II (FG-REB2)  
10X FastGene® FastCut Buffer (FG-REBHF)  
6X DNA Loading Buffer  
Sterile water

#### Recognition site



*For Research Use Only. Not for use in diagnostic procedures.*

ISO9001

**Source:** *Bacillus stearothermophilus* Y406

#### Reaction conditions

1X FastGene® Buffer II 60°C  
1X FastGene® FastCut Buffer, 60°C

#### FastGene® FastCut Buffer

FastGene® restriction enzyme can cut substrate DNA in 5-15 with FastGene® FastCut Buffer.

#### 1X FastGene® Buffer II

10 mM Tris-HCl (pH 7.9 at 25°C)  
50 mM NaCl  
10 mM MgCl<sub>2</sub>  
100 μg/ml BSA

#### Unit definition

One unit is defined as the amount of enzyme required for complete digestion of 1 μg bacteriophage λ at 60°C for 1 hr in 50 μl reaction mixtures.

#### Quality control

- Unit definition assay
- Overdigestion assay
- Endonuclease assay
- Extreme pure assay

**Dilution buffer:** FastGene® Diluent A

#### Heat Inactivation

BstY I can be inactivated at 80°C for 20 min.

#### Methylation sensitivity

*dam* methylation: Not sensitive  
*dcm* methylation: Not sensitive  
CpG methylation: Not sensitive

#### Prolonged incubation

A minimum amount of enzyme required to digest 1 μg substrate DNA for 16 hr; 0.13 U.

#### Relative activity in FastGene® Buffers

FastGene® Buffer I: 50%  
FastGene® Buffer II: 100%  
FastGene® Buffer III: 75%  
FastGene® Buffer IV: 100%  
FastGene® FastCut Buffer: 100%

#### Note

It is an isoschizomer of Xho II. It is not affected by *dam*, *dcm*, or mammalian CpG methylation.

#### Standard reaction condition

- Normal protocol

Component	Final Conc.	Volume
Substrate DNA	1 μg	X μl
10X FastGene® Buffer II	1 X	5 μl
BstY I	10 unit	1 μl
Sterile water		up to 50 μl

→ Incubate at 60°C for 1 hr

- Fast protocol

Component	Final Conc.	Volume
Substrate DNA	1 μg	X μl
10X FastGene® FastCut Buffer	1 X	5 μl
BstY I	10 unit	1 μl
Sterile water		up to 50 μl

→ Incubate at 60°C for 15 min

※ We recommend 5-10 units of enzyme per μg DNA and 10-20 units for genomic DNA in a 1 h digest.