



Restriction Enzyme

Nt.BstNB I




Cat.#	Size	Conc.
FG-NtBstNBI	1,000 units	10 units/µl

Store at -20°C

Supplied with: 10X FastGene® Buffer III (FG-REB3)
10X FastGene® FastCut Buffer (FG-REBHF)
6X DNA Loading Buffer
Sterile water

Recognition site



For Research Use Only. Not for use in diagnostic procedures.



Dilution buffer:

FastGene® Diluent A

Heat Inactivation

Nt.BstNB I can be inactivated at 80°C for 20 min.

Methylation sensitivity

dam methylation: Not sensitive

dcm methylation: Not sensitive

CpG methylation: Not sensitive

Prolonged incubation

A minimum amount of enzyme required to digest 1 µg substrate DNA for 16 hr; 0.5U.

Relative activity in FastGene® Buffers

FastGene® Buffer I:	0%
FastGene® Buffer II:	10%
FastGene® Buffer III:	100%
FastGene® Buffer IV:	0%
FastGene® FastCut Buffer:	100%

Note

Nt.BstNB I is a single cutter and cuts only the upper strand.

Source: *Bacillus stercorophilus* 33M

Reaction conditions

1X FastGene® Buffer III 55°C

1X FastGene® FastCut Buffer, 55°C

FastGene® FastCut Buffer

FastGene® restriction enzyme can cut substrate DNA in 5-15 with FastGene® FastCut Buffer.

1X FastGene® Buffer III

50 mM Tris-HCl (pH 7.9 at 25°C)

100 mM NaCl

10 mM MgCl₂

100 µg/ml BSA

Unit definition

One unit is defined as the amount of enzyme required for complete digestion of 1 µg pUC19 at 55°C for 1 hr in 50 µl reaction mixtures.

Quality control

- Unit definition assay
- Overdigestion assay
- Endonuclease assay
- Extreme pure assay

Standard reaction condition

- Normal protocol

Component	Final Conc.	Volume
Substrate DNA	1 µg	X µl
10X FastGene® Buffer III	1 X	5 µl
Nt.BstNB I	10 unit	1 µl
Sterile water		up to 50 µl

→ Incubate at 55°C for 1 hr

- Fast protocol

Component	Final Conc.	Volume
Substrate DNA	1 µg	X µl
10X FastGene® FastCut Buffer	1 X	5 µl
Nt.BstNB I	10 unit	1 µl
Sterile water		up to 50 µl

→ Incubate at 55°C for 15 min

※ We recommend 5-10 units of enzyme per µg DNA and 10-20 units for genomic DNA in a 1 h digest.